



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TNFRSF18 (Human) IP-WB Antibody Pair

Catalog # : H00008784-PW2

規格 : [1 Set]


List All

Specification

Product Description: This IP-WB antibody pair set comes with one antibody for immunoprecipitation and another to detect the precipitated protein in western blot.

Reactivity: Human

Quality Control Testing: Immunoprecipitation-Western Blot (IP-WB)



Immunoprecipitation of TNFRSF18 transfected lysate using rabbit polyclonal anti-TNFRSF18 and Protein A Magnetic Bead (U0007), and immunoblotted with mouse purified polyclonal anti-TNFRSF18.

Supplied Product: Antibody pair set content:
 1. Antibody pair for IP: rabbit polyclonal anti-TNFRSF18 (300 ul)
 2. Antibody pair for WB: mouse purified polyclonal anti-TNFRSF18 (50 ug)

Storage Instruction: Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -20°C storage immediately after use.

Applications

Immunoprecipitation-Western Blot

 Protocol Download

Gene Information

Entrez GeneID: [8784](#)

Gene Name: TNFRSF18

Gene Alias: AITR,GITR,GITR-D

Gene Description: tumor necrosis factor receptor superfamily, member 18

Omim ID: [603905](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor has been shown to have increased expression upon T-cell activation, and it is thought to play a key role in dominant immunological self-tolerance maintained by CD25(+)CD4(+) regulatory T cells. Knockout studies in mice also suggest the role of this receptor is in the regulation of CD3-driven T-cell activation and programmed cell death. Three alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported. [provided by RefSeq]

Other Designations: OTTHUMP00000001519, OTTHUMP00000001520, OTTHUMP00000001521, TNF receptor superfamily activation-inducible protein, activation-inducible TNFR family receptor, glucocorticoid-induced TNFR-related protein

Gene Pathway

[Cytokine-cytokine receptor interaction](#)

Related Disease

[Asthma](#) [Birth Weight](#) [Genetic Predisposition to Disease](#) [Glioblastoma](#) [Glioma](#) [Hepatitis C](#) [Hodgkin Disease](#) [Leukemia](#) [Leukemia, Myeloid, Acute](#) [Lymphoproliferative Disorders](#) [Meningeal Neoplasms](#) [Meningioma](#) [Waldenstrom Macroglobulinemia](#) [Werner syndrome](#)

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.