



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

NR1I2 (Human) IP-WB Antibody Pair

Catalog # : H00008856-PW2

規格 : [1 Set]


[List All](#)

Specification

Product Description: This IP-WB antibody pair set comes with one antibody for immunoprecipitation and another to detect the precipitated protein in western blot.

Reactivity: Human

Quality Control Testing: Immunoprecipitation-Western Blot (IP-WB)



Immunoprecipitation of NR1I2 transfected lysate using rabbit polyclonal anti-NR1I2 and Protein A Magnetic Bead ([U0007](#)), and immunoblotted with mouse purified polyclonal anti-NR1I2.

Supplied Product: Antibody pair set content:
 1. Antibody pair for IP: rabbit polyclonal anti-NR1I2 (300 ul)
 2. Antibody pair for WB: mouse purified polyclonal anti-NR1I2 (50 ug)

Storage Instruction: Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -20°C storage immediately after use.

Applications

Immunoprecipitation-Western Blot

 [Protocol Download](#)

Gene Information

Entrez GeneID: [8856](#)

Gene Name: NR1I2

Gene Alias: BXR, ONR1, PAR, PAR1, PAR2, PARq, PRR, PXR, SAR, SXR

Gene Description: nuclear receptor subfamily 1, group I, member 2

Omim ID: [603065](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: This gene product belongs to the nuclear receptor superfamily,

members of which are transcription factors characterized by a ligand-binding domain and a DNA-binding domain. The encoded protein is a transcriptional regulator of the cytochrome P450 gene CYP3A4, binding to the response element of the CYP3A4 promoter as a heterodimer with the 9-cis retinoic acid receptor RXR. It is activated by a range of compounds that induce CYP3A4, including dexamethasone and rifampicin. Several alternatively spliced transcripts encoding different isoforms, some of which use non-AUG (CUG) translation initiation codon, have been described for this gene. Additional transcript variants exist, however, they have not been fully characterized. [provided by RefSeq]

Other Designations: orphan nuclear receptor PXR, steroid and xenobiotic receptor

Related Disease

[Anemia](#) [Asthma](#) [Asthma](#) [Birth Weight](#) [Bone Neoplasms](#) [Breast cancer](#) [Breast Neoplasms](#) [Carcinoma, Non-Small-Cell Lung](#) [Cardiovascular Diseases](#) [Cholangitis, Sclerosing](#) [Cholestasis, Intrahepatic](#) [Colitis, Ulcerative](#) [Colon cancer](#) [Colonic Neoplasms](#) [Coronary Artery Disease](#) [Crohn Disease](#) [Crohn's disease](#) [Diabetes Mellitus, Type 2](#) [Drug-Induced Liver Injury](#)

... see more

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.