

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com



FAF6L (Hum a	an) IP-WB Antibody Pair	
Catalog # : H0001	0629-PW2 規格:[1 Set]	
List All		
Specification		Application Image
Product Description:	This IP-WB antibody pair set comes with one antibody for immunoprecipitation and another to detect the precipitated protein in western blot.	Immunoprecipitation-Western Blot
Reactivity:	Human	
Quality Control Testing:	Immunoprecipitation-Western Blot (IP-WB)	
Supplied Product:	Antibody pair set content: 1. Antibody pair for IP: rabbit polyclonal anti-TAF6L (300 ul) 2. Antibody pair for WB: mouse purified polyclonal anti-TAF6L (50 ug)	
Storage Instruction:	Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to - 20°C storage immediately after use.	
MSDS:	Download	
Applications		
Immunoprecipit	ation-Western Blot	
Gene Informatio	on	
Entrez GeneID:	10629	
Gene Name:	TAF6L	
Gene Alias:	FLJ11136,MGC4288,PAF65A	
Gene Description:	TAF6-like RNA polymerase II, p300/CBP-associated factor (PCAF)- associated factor, 65kDa	
Omim ID:	602946	
Gene Ontology	Hyperlink Degra 1 of 2	2010

Gene Summary:	Initiation of transcription by RNA polymerase II requires the activities of more than 70 polypeptides. The protein that coordinates these activities is transcription factor IID (TFIID), which binds to the core promoter to
	position the polymerase properly, serves as the scaffold for assembly of the remainder of the transcription complex, and acts as a channel for regulatory signals. TFIID is composed of the TATA-binding protein (TBP) and a group of evolutionarily conserved proteins known as TBP- associated factors or TAFs. TAFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modify general transcription factors (GTFs) to facilitate complex assembly and transcription initiation. This gene encodes a protein that is a component of the PCAF histone acetylase complex and structurally similar to one of the histone-like TAFs, TAF6. The PCAF histone acetylase complex, which is composed of more than 20 polypeptides some of which are TAFs, is required for myogenic transcription and differentiation. [provided by RefSeq
Other Designations:	TAF6-like RNA polymerase II,TAF6-like RNA polymerase II, p300/CBP- associated factor (PCAF)-associated factor, 65 kD,p300/CBP- associated factor (PCAF)-associated factor 65
Gene Pathway	

服務條款 | 隱私權政策 | 著作及商標 | 網站地圖 ©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.