



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

RBPMs monoclonal antibody (MF03), clone 1C12 (FITC)

Catalog Number: H00011030-MF03

Regulation Status: For research use only (RUO)

Product Description: FITC conjugated mouse monoclonal antibody raised against a full-length recombinant RBPMs.

Clone Name: 1C12

Immunogen: RBPMs (AAH03608, 1 a.a. ~ 219 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Sequence:

MNNGGKAEKENTPSEANLQEEVRTLFSGLPLDIKP
RELYLLFRPFKGYEGSLIKLTSKQPVGFVSFDSRSEAE
AAKNALNGIRFDPEIPQTLRLEFAKANTKMAKNKLVGT
PNPSTPLPNTVPQFIAREPYELTVPALYPSSPEVWAPY
PLYPAELAPALPPPAFTYPASLHAQCFSPEAKPNTVPV
CPLLQQIRFVSGNVFVTYQPTADQQRELPC

Host: Mouse

Interspecies Antigen Sequence: Mouse (99); Rat (99)

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product
page for detailed protocols

Conjugation: FITC

Isotype: IgG2a Kappa

Storage Buffer: In 1x PBS, pH 7.4

Storage Instruction: Store at -20°C or lower. Aliquot to
avoid repeated freezing and thawing.

Entrez GeneID: 11030

Gene Symbol: RBPMs

Gene Alias: HERMES

Gene Summary: This gene encodes a member of the
RRM family of RNA-binding proteins. The RRM domain

is between 80-100 amino acids in length and family members contain one to four copies of the domain. The RRM domain consists of two short stretches of conserved sequence called RNP1 and RNP2, as well as a few highly conserved hydrophobic residues. The protein encoded by this gene has a single, putative RRM domain in its N-terminus. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq]