



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

TARDBP monoclonal antibody (MA01J), clone 2E2-D3 (APC)

Catalog Number: H00023435-MA01J

Regulation Status: For research use only (RUO)

Product Description: APC conjugated mouse monoclonal antibody raised against a partial recombinant TARDBP.

This product is belong to Cell Culture Grade Antibody (CX Grade).

Clone Name: 2E2-D3

Immunogen: TARDBP (NP_031401.1, 1 a.a. ~ 260 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Sequence:

MSEYIRVTEDENDIEIPSEDDGTVLLSTVTAQFPGA
CGLRYRNPVSQCMRGVRLVEGILHAPDAGWGNLVYV
VNYPKDNKRKMDDETASSAVKVKRAVQKTSIDLVLGL
PWKTTEQDLKEYFSTFGEVLMVQVKKDLKTGHSGKF
GFVRFTEYETQVKVMSQRHMIDGRWCCKLPNSKQS
QDEPLRSRKVFVGRCTEDMTEDELREFFSQYGDVMD
VFIPKPFRAFAFVTFADDQIAQSLCGEDLIKGISVHISN
A

Host: Mouse

Interspecies Antigen Sequence: Mouse (96)

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product
page for detailed protocols

Conjugation: APC

Preparation Method: Cell Culture Production

Isotype: IgG1 Kappa

Storage Buffer: In 1x PBS, pH 7.4

Storage Instruction: Store at -20°C or lower. Aliquot to
avoid repeated freezing and thawing.

Entrez GeneID: 23435

Gene Symbol: TARDBP

Gene Alias: ALS10, TDP-43

Gene Summary: HIV-1, the causative agent of acquired immunodeficiency syndrome (AIDS), contains an RNA genome that produces a chromosomally integrated DNA during the replicative cycle. Activation of HIV-1 gene expression by the transactivator Tat is dependent on an RNA regulatory element (TAR) located downstream of the transcription initiation site. The protein encoded by this gene is a transcriptional repressor that binds to chromosomally integrated TAR DNA and represses HIV-1 transcription. In addition, this protein regulates alternate splicing of the CFTR gene. A similar pseudogene is present on chromosome 20. [provided by RefSeq]