



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

Anti-Tenecteplase (Human) ELISA Kit

Catalog Number: KA7149

Regulation Status: For research use only (RUO)

Product Description: Anti-Tenecteplase (Human)
ELISA Kit is a kit used for the quantitative measurement of antibodies to Tenecteplase in serum and plasma.

Calibration Range: 0 to 8,000 ng/mL

Limit of Detection: 28 ng/mL

Suitable Sample: Citrate Plasma, EDTA Plasma, Heparin Plasma, Serum

Sample Volume: 100 μ L

Detection Method: Colorimetric

Absorbance (nm): 450

Assay type: Quantitative

Spiking Recovery: 80%-120%

Regulation Status: For research use only (RUO)

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Storage Instruction: Store the kit at 4°C.

Entrez GeneID: 5327

Gene Symbol: PLAT

Gene Alias: DKFZp686I03148, T-PA, TPA

Gene Summary: This gene encodes tissue-type plasminogen activator, a secreted serine protease which converts the proenzyme plasminogen to plasmin, a fibrinolytic enzyme. Tissue-type plasminogen activator is synthesized as a single chain which is cleaved by plasmin to a two chain disulfide linked protein. This enzyme plays a role in cell migration and tissue remodeling. Increased enzymatic activity causes

hyperfibrinolysis, which manifests as excessive bleeding; decreased activity leads to hypofibrinolysis which can result in thrombosis or embolism. Alternative splicing of this gene results in multiple transcript variants encoding different isoforms. [provided by RefSeq]