



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

SELP monoclonal antibody, clone 4AW12 (PE)

Catalog Number: MAB12413

Regulatory Status: For research use only (RUO)

Product Description: Mouse monoclonal antibody raised against human SELP.

Clone Name: 4AW12

Immunogen: SELP

Host: Mouse

Reactivity: Human

Applications: Flow Cyt, IF
(See our web site product page for detailed applications information)

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Conjugation: PE

Purification: Protein A/G affinity chromatography

Isotype: IgG1

Recommend Usage: Flow cytometry
Immunofluorescence
The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS, pH 7.4 (0.09% sodium azide, 0.2% BSA)

Storage Instruction: Store at 4°C. Do not freeze.

Entrez GeneID: 6403

Gene Symbol: SELP

Gene Alias: CD62, CD62P, FLJ45155, GMP140, GRMP, LECAM3, PADGEM, PSEL

Gene Summary: This gene encodes a 140 kDa protein that is stored in the alpha-granules of platelets and Weibel-Palade bodies of endothelial cells. This protein redistributes to the plasma membrane during platelet activation and degranulation and mediates the interaction of activated endothelial cells or platelets with leukocytes. The membrane protein is a calcium-dependent receptor that binds to sialylated forms of Lewis blood group carbohydrate antigens on neutrophils and monocytes. Alternative splice variants may occur but are not well documented. [provided by RefSeq]