



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

HMGCR monoclonal antibody, clone CL0259

Catalog Number: MAB15579

Regulatory Status: For research use only (RUO)

Product Description: Mouse monoclonal antibody raised against partial recombinant human HMGCR.

Clone Name: CL0259

Immunogen: Recombinant protein corresponding to human HMGCR.

Sequence:

HIPAYKLETLMETHERGVSI RRQLLSKKLSEPSSLQYLP
YRDYNYSLVMGACCENVIGYMPIVGVAGPLCLDEKE
FQVPMATTEGCLVA

Host: Mouse

Epitope: This antibody binds to an epitope located within the peptide sequence TLMETHERGVSI RRQ as determined by overlapping synthetic peptides.

Reactivity: Human

Applications: IHC-P, WB-Ce

(See our web site product page for detailed applications information)

Protocols: See our web site at

<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Purification: Protein A purification

Isotype: IgG2a

Recommend Usage: Immunohistochemistry

(Formalin/PFA-fixed paraffin-embedded sections)

(1:50-1:200)

Western Blot (1:500-1:1000)

The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS, pH 7.2 (40% glycerol, 0.02%

sodium azide).

Storage Instruction: Store at 4°C. For long term storage store at -20°C.

Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 3156

Gene Symbol: HMGCR

Gene Alias: -

Gene Summary: HMG-CoA reductase is the rate-limiting enzyme for cholesterol synthesis and is regulated via a negative feedback mechanism mediated by sterols and non-sterol metabolites derived from mevalonate, the product of the reaction catalyzed by reductase. Normally in mammalian cells this enzyme is suppressed by cholesterol derived from the internalization and degradation of low density lipoprotein (LDL) via the LDL receptor. Competitive inhibitors of the reductase induce the expression of LDL receptors in the liver, which in turn increases the catabolism of plasma LDL and lowers the plasma concentration of cholesterol, an important determinant of atherosclerosis. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]