



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

FBLN1 monoclonal antibody, clone CL0337

Catalog Number: MAB15607

Regulatory Status: For research use only (RUO)

Product Description: Mouse monoclonal antibody raised against partial recombinant human FBLN1.

Clone Name: CL0337

Immunogen: Recombinant protein corresponding to human FBLN1.

Sequence:

CESGIHNCPLPDFICQNTLGSRFCRKLQCKSGFIQDAL
GNCIDINECLISAPCPIGHTCINTEGSYTCQKNVPNCG
RGYHLNNEEGTRCVDVDECAPPAEPCGKGHRCVNSPG
SFRCECKTGYFYFDGISRMCVDVNE

Host: Mouse

Reactivity: Human

Applications: IF, IHC-P, WB-Ti

(See our web site product page for detailed applications information)

Protocols: See our web site at

<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Purification: Protein A purification

Isotype: IgG1

Recommend Usage: Immunofluorescence (1-4 ug/mL)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (1:50-1:200)

Western Blot (1:500-1:1000)

The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS, pH 7.2 (40% glycerol, 0.02% sodium azide).

Storage Instruction: Store at 4°C. For long term

storage store at -20°C.

Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 2192

Gene Symbol: FBLN1

Gene Alias: FBLN

Gene Summary: Fibulin 1 is a secreted glycoprotein that becomes incorporated into a fibrillar extracellular matrix. Calcium-binding is apparently required to mediate its binding to laminin and nidogen. It mediates platelet adhesion via binding fibrinogen. Four splice variants which differ in the 3' end have been identified. Each variant encodes a different isoform, but no functional distinctions have been identified among the four variants. [provided by RefSeq]