



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

SDHB monoclonal antibody, clone CL0346

Catalog Number: MAB15611

Regulatory Status: For research use only (RUO)

Product Description: Mouse monoclonal antibody raised against partial recombinant human SDHB.

Clone Name: CL0346

Immunogen: Recombinant protein corresponding to human SDHB.

Sequence:

EGKQQYLQSIEREKLDGLYECILCACCTSCPSYWW
NGDKYLGPAVLMQAYRWMIDSRDDFTEERLAKLQDP
FSLYRCHTIMNCTRTPKGLNPGKAIAEIKKMMATY

Host: Mouse

Reactivity: Human

Applications: IHC-P, WB-Ce

(See our web site product page for detailed applications information)

Protocols: See our web site at

<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Purification: Protein A purification

Isotype: IgG2a

Recommend Usage: Immunohistochemistry

(Formalin/PFA-fixed paraffin-embedded sections)
(1:200-1:500)

Western Blot (1:500-1:1000)

The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS, pH 7.2 (40% glycerol, 0.02% sodium azide).

Storage Instruction: Store at 4°C. For long term storage store at -20°C.

Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 6390

Gene Symbol: SDHB

Gene Alias: FLJ92337, IP, PGL4, SDH, SDH1, SDHIP

Gene Summary: Complex II of the respiratory chain, which is specifically involved in the oxidation of succinate, carries electrons from FADH to CoQ. The complex is composed of four nuclear-encoded subunits and is localized in the mitochondrial inner membrane. The iron-sulfur subunit is highly conserved and contains three cysteine-rich clusters which may comprise the iron-sulfur centers of the enzyme. Sporadic and familial mutations in this gene result in paragangliomas and pheochromocytoma, and support a link between mitochondrial dysfunction and tumorigenesis. [provided by RefSeq]