



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

MKI67IP monoclonal antibody, clone CL2237

Catalog Number: MAB15733

Regulatory Status: For research use only (RUO)

Product Description: Mouse monoclonal antibody raised against partial recombinant human MKI67IP.

Clone Name: CL2237

Immunogen: Recombinant protein corresponding to human MKI67IP.

Sequence:

IDYDFPSLILQKTESISKTNRQTSTKGQVLRKKKKKVS
GTLDTPEKTVDSQGPTPVCTPTFLERRKSQVAELNDDD
KDDEI

Host: Mouse

Epitope: This antibody binds to an epitope located within the peptide sequence IDYDFPSLILQKTES as determined by overlapping synthetic peptides.

Reactivity: Human

Applications: IHC-P, WB-Ce

(See our web site product page for detailed applications information)

Protocols: See our web site at

<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Purification: Protein A purification

Isotype: IgG1

Recommend Usage: Immunohistochemistry

(Formalin/PFA-fixed paraffin-embedded sections)
(1:2500-1:5000)

Western Blot (1:500-1:1000)

The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS, pH 7.2 (40% glycerol, 0.02%

sodium azide).

Storage Instruction: Store at 4°C. For long term storage store at -20°C.

Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 84365

Gene Symbol: MKI67IP

Gene Alias: NIFK, Nopp34

Gene Summary: This gene encodes a protein that interacts with the forkhead-associated domain of the Ki-67 antigen. The encoded protein may bind RNA and may play a role in mitosis and cell cycle progression. Multiple pseudogenes exist on chromosomes 5, 10, 12, 15, and 19.[provided by RefSeq]