



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

ASGPR1 monoclonal antibody (FITC)

Catalog Number: MAB23669-M01

Regulation Status: For research use only (RUO)

Product Description: FITC conjugated ASGPR1 mouse monoclonal antibody.

Immunogen: Human recombinant ASGPR1

Host: Mouse

Amenable Platform: [LiquidCell™ negative enrichment cell isolation](#)

Regulatory Status: Research Use Only

Instrument Compatibility: Ventana™ BenchMark Stainer

Available Test: 16 assays

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Conjugation: FITC

Concentration: 0.5 mg/mL

Isotype: IgG1, kappa

Recommend Usage: Immunofluorescence (10 ug/mL)
LiquidCell™ SuperSlide™ (6 uL per slide)

Storage Buffer: In 1x PBS, pH 7.4

Storage Instruction: Store at 2-8°C. Do not freeze.

Entrez GeneID: 432

Gene Symbol: ASGR1

Gene Alias: HL-1, ASGPR, ASGPR1, CLEC4H1

Gene Summary: Partially deglycosylated plasma glycoproteins and immunoglobulin IgA2 allotypes are

efficiently and specifically removed from circulation by a receptor-mediated process. The asialoglycoprotein receptor binds to desialylated (galactosyl-terminal) glycoproteins. It transports these glycoproteins via a series of membrane vesicles and tubules to an acidic-sorting organelle where the receptor and ligand dissociate. Then the receptor is recycled back to the cell surface and the ligand is transported to the lysosomes for degradation. [provided by RefSeq]