



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

ITGAL monoclonal antibody, clone TP1/40

Catalog Number: MAB4965

Regulatory Status: For research use only (RUO)

Product Description: Mouse monoclonal antibody raised against native ITGAL.

Clone Name: TP1/40

Immunogen: PMA and ionomycin activated human T cells blast.

Host: Mouse

Reactivity: Human

Applications: Flow Cyt, IF
(See our web site product page for detailed applications information)

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Purification: Protein A/G purification

Isotype: IgG1

Recommend Usage: Flow Cytometry (20 μ L/ 10^6 cells)
The optimal working dilution should be determined by the end user.

Storage Buffer: In buffer containing 1% BSA, pH 7.2 (0.09% sodium azide).

Storage Instruction: Store in the dark at 4°C. Avoid prolonged exposure to light.

Entrez GeneID: 3683

Gene Symbol: ITGAL

Gene Alias: CD11A, LFA-1, LFA1A

Gene Summary: ITGAL encodes the integrin alpha L

chain. Integrins are heterodimeric integral membrane proteins composed of an alpha chain and a beta chain. This I-domain containing alpha integrin combines with the beta 2 chain (ITGB2) to form the integrin lymphocyte function-associated antigen-1 (LFA-1), which is expressed on all leukocytes. LFA-1 plays a central role in leukocyte intercellular adhesion through interactions with its ligands, ICAMs 1-3 (intercellular adhesion molecules 1 through 3), and also functions in lymphocyte costimulatory signaling. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]