



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



Datasheet

VHH-His tag NanoAb™ Targeting Human CD30, clone 9063

Catalog Number: NAB00025-M01J

Regulation Status: For research use only (RUO)

Product Description: VHH-His tag NanoAb™ targeting human CD30 native protein.

Clone Name: 9063

Immunogen: Human CD30 recombinant protein

Sequence: Available for licensing

Antibody Species: Camelid

Epitope: Monospecific

Tag: 6x-His Tag at C-terminus

Conjugation: Unconjugated

Affinity: Not measured

Form: Liquid

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product
page for detailed protocols

Conjugation: Unconjugated

Preparation Method: Mammalian cell (HEK293)
expression system

Purification: Ni-IDA (Iminodiacetic acid) resin

Recommend Usage: Flow cytometry
ELISA

The optimal working dilution should be determined by
the end user.

Storage Buffer: In PBS, pH 7.4

Storage Instruction: Store at -80°C.
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 943

Gene Symbol: TNFRSF8

Gene Alias: CD30, D1S166E, KI-1

Gene Summary: The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is expressed by activated, but not by resting, T and B cells. TRAF2 and TRAF5 can interact with this receptor, and mediate the signal transduction that leads to the activation of NF-kappaB. This receptor is a positive regulator of apoptosis, and also has been shown to limit the proliferative potential of autoreactive CD8 effector T cells and protect the body against autoimmunity. Two alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported. [provided by RefSeq]