



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

VHH-His tag NanoAb™ Targeting Human GM-CSF, clone P151

Catalog Number: NAB00082-M01J

Regulation Status: For research use only (RUO)

Product Description: VHH-His tag NanoAb™ targeting human GM-CSF native protein.

Clone Name: P151

Immunogen: Human GM-CSF recombinant protein

Sequence: Available for licensing

Antibody Species: Camelid

Epitope: Monospecific

Tag: 6x-His Tag at C-terminus

Conjugation: Unconjugated

Affinity: Not measured

Form: Liquid

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product
page for detailed protocols

Conjugation: Unconjugated

Preparation Method: Mammalian cell (HEK293)
expression system

Purification: Ni-IDA (Iminodiacetic acid) resin

Recommend Usage: Flow cytometry
ELISA

The optimal working dilution should be determined by
the end user.

Storage Buffer: In PBS, pH 7.4

Storage Instruction: Store at -80°C.
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 1437

Gene Symbol: CSF2

Gene Alias: GMCSF, MGC131935, MGC138897

Gene Summary: The protein encoded by this gene is a cytokine that controls the production, differentiation, and function of granulocytes and macrophages. The active form of the protein is found extracellularly as a homodimer. This gene has been localized to a cluster of related genes at chromosome region 5q31, which is known to be associated with interstitial deletions in the 5q- syndrome and acute myelogenous leukemia. Other genes in the cluster include those encoding interleukins 4, 5, and 13. [provided by RefSeq]