



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

FAS (Human) Recombinant Protein

Catalog Number: P10027

Regulation Status: For research use only (RUO)

Product Description: Human FAS (P25445-1, Gln26-Asn173) partial recombinant protein with hFc tag at C-terminus expressed in HEK293 cells.

Sequence: Gln26-Asn173

Host: Human

Theoretical MW (kDa): 43.4

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Lyophilized

Preparation Method: Mammalian cell (HEK293) expression system

Purity: > 95% as determined by Tris-Bis PAGE; > 95% as determined by HPLC

Endotoxin Level: < 1 EU per 1 ug of protein (determined by LAL method)

Recommend Usage: Biological Activity

ELISA

SEC-HPLC

Tris-Bis PAGE

The optimal working dilution should be determined by the end user.

Storage Buffer: Lyophilized from filtered solution in PBS, pH 7.4 (5% trehalose).

Storage Instruction: After reconstitution with deionized water to a final concentration more than 100 ug/ml, store at 4°C for 1 week. For long term storage, store at -80°C for 1 year.

Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 355

Gene Symbol: FAS

Gene Alias: ALPS1A, APO-1, APT1, CD95, FAS1, FASTM, TNFRSF6

Gene Summary: The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor contains a death domain. It has been shown to play a central role in the physiological regulation of programmed cell death, and has been implicated in the pathogenesis of various malignancies and diseases of the immune system. The interaction of this receptor with its ligand allows the formation of a death-inducing signaling complex that includes Fas-associated death domain protein (FADD), caspase 8, and caspase 10. The autoproteolytic processing of the caspases in the complex triggers a downstream caspase cascade, and leads to apoptosis. This receptor has been also shown to activate NF-kappaB, MAPK3/ERK1, and MAPK8/JNK, and is found to be involved in transducing the proliferating signals in normal diploid fibroblast and T cells. At least eight alternatively spliced transcript variants have been described, some of which are candidates for nonsense-mediated decay (NMD). The isoforms lacking the transmembrane domain may negatively regulate the apoptosis mediated by the full length isoform. [provided by RefSeq]