

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

Datasheet

KLRK1 (Human) Recombinant Protein

Catalog Number: P10035

Regulation Status: For research use only (RUO)

Product Description: Human KLRK1 (P26718,

Phe78-Val216) partial recombinant protein with hFc-Flag

tag at N-terminus expressed in HEK293 cells.

Sequence: Phe78-Val216

Host: Human

Theoretical MW (kDa): 43.4

Protocols: See our web site at

http://www.abnova.com/support/protocols.asp or product

page for detailed protocols

Form: Lyophilized

Preparation Method: Mammalian cell (HEK293)

expression system

Purity: > 95% as determined by Tris-Bis PAGE;> 95%

as determined by HPLC

Endotoxin Level: < 1 EU per 1 ug of protein

(determined by LAL method)

Recommend Usage: Biological Activity

ELISA SEC-HPLC Tris-Bis PAGE

The optimal working dilution should be determined by

the end user.

Storage Buffer: Lyophilized from filtered solution in

PBS, pH 7.4 (5% trehalose).

Storage Instruction: After reconstitution with deionized water to a final concentration more than 100 ug/ml, store at 4°C for 1 week. For long term storage, store at -80°C

for 1 year.

Aliquot to avoid repeated freezing and thawing.

Entrez GenelD: 22914

Gene Symbol: KLRK1

Gene Alias: CD314, D12S2489E, FLJ17759, FLJ75772,

KLR, NKG2-D, NKG2D

Gene Summary: Natural killer (NK) cells are

lymphocytes that can mediate lysis of certain tumor cells and virus-infected cells without previous activation. They can also regulate specific humoral and cell-mediated immunity. NK cells preferentially express several calcium-dependent (C-type) lectins, which have been implicated in the regulation of NK cell function. This gene encodes a member of the NKG2 family, and the encoded transmembrane protein is characterized by a type II membrane orientation (extracellular C terminus) and the presence of a C-type lectin domain. The NKG2 gene family is located within the NK complex, a region that contains several C-type lectin genes preferentially

expressed in NK cells. [provided by RefSeq]