



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

TNFRSF18 (Human) Recombinant Protein

Catalog Number: P10128

Regulation Status: For research use only (RUO)

Product Description: Human TNFRSF18 (Q9Y5U5-1, Gln26-Glu161) partial recombinant protein with His tag at C-terminus expressed in HEK293 cells.

Sequence: Gln26-Glu161

Host: Human

Theoretical MW (kDa): 15.4

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Lyophilized

Preparation Method: Mammalian cell (HEK293) expression system

Purity: > 95% as determined by Tris-Bis PAGE; > 95% as determined by HPLC

Endotoxin Level: < 1 EU per 1 ug of protein (determined by LAL method)

Recommend Usage: Biological Activity

ELISA

SEC-HPLC

SPR

Tris-Bis PAGE

The optimal working dilution should be determined by the end user.

Storage Buffer: Lyophilized from filtered solution in PBS, pH 7.4 (8% trehalose).

Storage Instruction: After reconstitution with deionized water to a final concentration more than 100 ug/ml, store at 4°C for 1 week. For long term storage, store at -80°C for 1 year.

Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 8784

Gene Symbol: TNFRSF18

Gene Alias: AITR, GITR, GITR-D

Gene Summary: The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor has been shown to have increased expression upon T-cell activation, and it is thought to play a key role in dominant immunological self-tolerance maintained by CD25(+)CD4(+) regulatory T cells. Knockout studies in mice also suggest the role of this receptor is in the regulation of CD3-driven T-cell activation and programmed cell death. Three alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported. [provided by RefSeq]