

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

Datasheet

ERBB2 (Human) Recombinant Protein

Catalog Number: P10133

Regulation Status: For research use only (RUO)

Product Description: Human ERBB2 (P04626-1, Thr23-Thr652) partial recombinant protein with His-Avi tag at C-terminus expressed in HEK293 cells.

Sequence: Thr23-Thr652

Host: Human

Theoretical MW (kDa): 72.3

Protocols: See our web site at

http://www.abnova.com/support/protocols.asp or product

page for detailed protocols

Form: Lyophilized

Preparation Method: Mammalian cell (HEK293)

expression system

Purity: > 95% as determined by Tris-Bis PAGE;> 95%

as determined by HPLC

Endotoxin Level: < 1 EU per 1 ug of protein

(determined by LAL method)

Recommend Usage: Biological Activity

ELISA SEC-HPLC Tris-Bis PAGE

The optimal working dilution should be determined by

the end user.

Storage Buffer: Lyophilized from filtered solution in

PBS, pH 7.4 (5% trehalose).

Storage Instruction: After reconstitution with deionized water to a final concentration more than 100 ug/ml, store at 4°C for 1 week. For long term storage, store at -80°C

for 1 year.

Aliquot to avoid repeated freezing and thawing.

Entrez GenelD: 2064

Gene Symbol: ERBB2

Gene Alias: CD340, HER-2, HER-2/neu, HER2, NEU,

NGL, TKR1

Gene Summary: This gene encodes a member of the epidermal growth factor (EGF) receptor family of receptor tyrosine kinases. This protein has no ligand binding domain of its own and therefore cannot bind growth factors. However, it does bind tightly to other ligand-bound EGF receptor family members to form a heterodimer, stabilizing ligand binding and enhancing kinase-mediated activation of downstream signalling pathways, such as those involving mitogen-activated protein kinase and phosphatidylinositol-3 kinase. Allelic variations at amino acid positions 654 and 655 of isoform a (positions 624 and 625 of isoform b) have been reported, with the most common allele. Ile654/Ile655, shown here. Amplification and/or overexpression of this gene has been reported in numerous cancers, including breast and ovarian tumors. Alternative splicing results in several additional transcript variants, some encoding different isoforms and others that have not been fully characterized. [provided by RefSeq]