



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



Datasheet

TNFRSF17 (Human) Recombinant Protein

Catalog Number: P10237

Regulation Status: For research use only (RUO)

Product Description: Human TNFRSF17 (Q02223-1, Met1-Ala54) partial recombinant protein with His-Avi tag at C-terminus expressed in HEK293 cells.

Sequence: Met1-Ala54

Host: Human

Theoretical MW (kDa): 8.9

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Lyophilized

Preparation Method: Mammalian cell (HEK293) expression system

Purity: > 95% as determined by Tris-Bis PAGE; > 95% as determined by HPLC

Endotoxin Level: < 1 EU per 1 ug of protein (determined by LAL method)

Recommend Usage: Biological Activity

ELISA

SEC-HPLC

Tris-Bis PAGE

The optimal working dilution should be determined by the end user.

Storage Buffer: Lyophilized from filtered solution in PBS, pH 7.4 (5% trehalose).

Storage Instruction: After reconstitution with deionized water to a final concentration more than 100 ug/ml, store at 4°C for 1 week. For long term storage, store at -80°C for 1 year.

Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 608

Gene Symbol: TNFRSF17

Gene Alias: BCM, BCMA, CD269

Gene Summary: The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is preferentially expressed in mature B lymphocytes, and may be important for B cell development and autoimmune response. This receptor has been shown to specifically bind to the tumor necrosis factor (ligand) superfamily, member 13b (TNFSF13B/TALL-1/BAFF), and to lead to NF-kappaB and MAPK8/JNK activation. This receptor also binds to various TRAF family members, and thus may transduce signals for cell survival and proliferation. [provided by RefSeq]