

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

#### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

#### **Datasheet**

# CLEC12A (Human) Recombinant Protein

Catalog Number: P10293

Regulation Status: For research use only (RUO)

**Product Description:** Human CLEC12A (Q5QGZ9-2, His65-Ala265) partial recombinant protein with His tag at

N-terminus expressed in HEK293 cells.

Sequence: His65-Ala265

Host: Human

Theoretical MW (kDa): 24.6

Protocols: See our web site at

http://www.abnova.com/support/protocols.asp or product

page for detailed protocols

Form: Lyophilized

Preparation Method: Mammalian cell (HEK293)

expression system

**Purity:** > 95% as determined by Tris-Bis PAGE;> 95%

as determined by HPLC

Endotoxin Level: < 1 EU per 1 ug of protein

(determined by LAL method)

Recommend Usage: Biological Activity

ELISA

Tris-Bis PAGE

The optimal working dilution should be determined by

the end user.

Storage Buffer: Lyophilized from filtered solution in

PBS, pH 7.4 (5% trehalose).

**Storage Instruction:** After reconstitution with deionized water to a final concentration more than 100 ug/ml, store at 4°C for 1 week. For long term storage, store at -80°C

for 1 year.

Aliquot to avoid repeated freezing and thawing.

Entrez GenelD: 160364

Gene Symbol: CLEC12A

Gene Alias: CLL-1, CLL1, DCAL-2, MGC70602, MICL

Gene Summary: This gene encodes a member of the C-type lectin/C-type lectin-like domain (CTL/CTLD) superfamily. Members of this family share a common protein fold and have diverse functions, such as cell adhesion, cell-cell signalling, glycoprotein turnover, and roles in inflammation and immune response. The protein encoded by this gene is a negative regulator of granulocyte and monocyte function. Several alternatively spliced transcript variants of this gene have been described, but the full-length nature of some of these variants has not been determined. This gene is closely linked to other CTL/CTLD superfamily members in the natural killer gene complex region on chromosome 12p13. [provided by RefSeq]