



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

CSF1 (Human) Recombinant Protein

Catalog Number: P10308

Regulation Status: For research use only (RUO)

Product Description: Human CSF1 recombinant protein with His tag at the C-terminus expressed in HEK293 cells.

Sequence: Glu33-Gln181

Host: Human

Theoretical MW (kDa): 19.21

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Lyophilized

Preparation Method: Mammalian cell (HEK293) expression system

Purity: > 95% by SDS-PAGE

Endotoxin Level: < 0.1 EU per 1 ug as determined by the LAL method.

Recommend Usage: SDS-PAGE

The optimal working dilution should be determined by the end user.

Storage Buffer: Lyophilized from a 0.2 um filtered solution of PBS, pH 7.4.

Storage Instruction: Store at -20°C for 12 months in lyophilized state.
After reconstitution with deionized water, store at -20 or -80°C for 1 month.
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 1435

Gene Symbol: CSF1

Gene Alias: MCSF, MGC31930

Gene Summary: The protein encoded by this gene is a

cytokine that controls the production, differentiation, and function of macrophages. The active form of the protein is found extracellularly as a disulfide-linked homodimer, and is thought to be produced by proteolytic cleavage of membrane-bound precursors. The encoded protein may be involved in development of the placenta. Four transcript variants encoding three different isoforms have been found for this gene. [provided by RefSeq]