



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



Datasheet

CD40LG Trimer (Human) Recombinant Protein (Biotin)

Catalog Number: P10385

Regulation Status: For research use only (RUO)

Product Description: Human CD40LG Trimer (P29965, Met113-Leu261) partial recombinant protein with His-Flag tag at the N-Terminus expressed in HEK293 cells.

Sequence: Met113-Leu261

Host: Human

Theoretical MW (kDa): 52.4

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Lyophilized

Conjugation: Biotin

Preparation Method: Mammalian cell (HEK293) expression system

Purity: > 95% by Tris-Bis PAGE
> 95% by HPLC

Endotoxin Level: < 0.1 EU per 1 ug as determined by the LAL method.

Activity: The EC₅₀ was 0.70 ug/mL, measured by ELISA at 2 ug/mL.

Recommend Usage: Biological Activity

ELISA

Tris-Bis PAGE

SEC-HPLC

The optimal working dilution should be determined by the end user.

Storage Buffer: Lyophilized from 0.22 um filtered solution in PBS, pH 7.4. (8% trehalose).

Storage Instruction: Store at -20°C for 12 months. After reconstitution, store at 4°C for 2-7 days, or store at -80°C for 3-6 months.

Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 959

Gene Symbol: CD40LG

Gene Alias: CD154, CD40L, HIGM1, IGM, IMD3, T-BAM, TNFSF5, TRAP, gp39, hCD40L

Gene Summary: The protein encoded by this gene is expressed on the surface of T cells. It regulates B cell function by engaging CD40 on the B cell surface. A defect in this gene results in an inability to undergo immunoglobulin class switch and is associated with hyper-IgM syndrome. [provided by RefSeq]