SZABO SCANDIC

Part of Europa Biosite

Produktinformation

- Forschungsprodukte & Biochemikalien
- Zellkultur & Verbrauchsmaterial
- Diagnostik & molekulare Diagnostik
- Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!

Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic

9F, No. 108, Jhouzih St., Taipei, Taiwan
Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218
E-mail: sales@abnova.com

Datasheet

DLL3 (Human) Recombinant Protein (Biotin)

Catalog Number: P10393

Regulation Status: For research use only (RUO)

Product Description: Human DLL3 (Q9NYJ7-1, Ala27-Arg490) partial recombinant protein with His tag at the N-Terminus expressed in HEK293 cells.

Sequence: Ala27-Arg490

Host: Human

Theoretical MW (kDa): 50.4

Protocols: See our web site at http://www.abnova.com/support/protocols.asp or product page for detailed protocols

Form: Lyophilized

Conjugation: Biotin

Preparation Method: Mammalian cell (HEK293) expression system

Purity: > 95% by Tris-Bis PAGE

Endotoxin Level: < 0.1 EU per 1 ug as determined by the LAL method.

Activity: The EC_{50} was 17.5 ng/mL, measured by ELISA at 2 ug/mL.

Recommend Usage: Biological Activity
ELISA
Tris-Bis PAGE
The optimal working dilution should be determined by the end user.

Storage Buffer: Lyophilized from 0.22 um filtered solution in PBS, pH 7.4. (8% trehalose).

Storage Instruction: Store at -20°C for 12 months.
After reconstitution, store at 4°C for 2-7 days, or store at -80°C for 3-6 months.
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 10683

Gene Symbol: DLL3

Gene Alias: SCDO1

Gene Summary: This gene encodes a member of the delta protein ligand family. This family functions as Notch ligands that are characterized by a DSL domain, EGF repeats, and a transmembrane domain. Mutations in this gene cause autosomal recessive spondylocostal dysostosis 1. Two transcript variants encoding distinct isoforms have been identified for this gene. [provided by RefSeq]