



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

IL5 (Human) Recombinant Protein (Biotin)

Catalog Number: P10434

Regulation Status: For research use only (RUO)

Product Description: Human IL5 (P05113, Ile20-Ser134) partial recombinant protein with His-Avi tag at the N-Terminus expressed in HEK293 cells.

Sequence: Ile20-Ser134

Host: Human

Theoretical MW (kDa): 16.1

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Lyophilized

Conjugation: Biotin

Preparation Method: Mammalian cell (HEK293) expression system

Purity: > 95% by Tris-Bis PAGE
> 95% by HPLC

Endotoxin Level: < 0.1 EU per 1 ug as determined by the LAL method.

Activity: The EC_{50} was 68.2 ng/mL, measured by ELISA at 2 ug/mL. The affinity constant was 7.57 nM, determined by SPR assay.

Recommend Usage: Biological Activity

ELISA

Tris-Bis PAGE

SEC-HPLC

SPR

The optimal working dilution should be determined by the end user.

Storage Buffer: Lyophilized from 0.22 um filtered solution in PBS, pH 7.4. (8% trehalose).

Storage Instruction: Store at -20°C for 24 months.

After reconstitution, store at 4°C for 2-7 days, or store at -80°C for 3-6 months.

Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 3567

Gene Symbol: IL5

Gene Alias: EDF, IL-5, TRF

Gene Summary: The protein encoded by this gene is a cytokine that acts as a growth and differentiation factor for both B cells and eosinophils. This cytokine is a main regulator of eosinopoiesis, eosinophil maturation and activation. The elevated production of this cytokine is reported to be related to asthma or hypereosinophilic syndromes. The receptor of this cytokine is a heterodimer, whose beta subunit is shared with the receptors for interleukine 3 (IL3) and colony stimulating factor 2 (CSF2/GM-CSF). This gene, together with those for interleukin 4 (IL4), interleukin 13 (IL13), and CSF2, form a cytokine gene cluster on chromosome 5. This cytokine, IL4, and IL13 are found to be regulated coordinately by long-range regulatory elements spread over 120 kilobases on chromosome 5q31. [provided by RefSeq]