

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

Datasheet

KLRC1 (Human) Recombinant Protein (Biotin)

Catalog Number: P10474

Regulation Status: For research use only (RUO)

Product Description: Human KLRC1 (P26715-1, Arg100-Leu233) partial recombinant protein with His-Avi tag at the N-Terminus expressed in HEK293 cells.

Sequence: Arg100-Leu233

Host: Human

Theoretical MW (kDa): 18.3

Protocols: See our web site at

http://www.abnova.com/support/protocols.asp or product

page for detailed protocols

Form: Lyophilized

Conjugation: Biotin

Preparation Method: Mammalian cell (HEK293)

expression system

Purity: > 95% by Tris-Bis PAGE

Endotoxin Level: < 0.1 EU per 1 ug as determined by

the LAL method.

Activity: The EC₅₀ was 5.3 ng/mL, measured by ELISA

at 1 ug/mL.

Recommend Usage: Biological Activity

ELISA

Tris-Bis PAGE

The optimal working dilution should be determined by

the end user.

Storage Buffer: Lyophilized from 0.22 um filtered

solution in PBS, pH 7.4. (8% trehalose).

Storage Instruction: Store at -20°C for 12 months. After reconstitution, store at 4°C for 2-7 days, or store at

-80°C for 3-6 months.

Aliquot to avoid repeated freezing and thawing.

Entrez GenelD: 3821

Gene Symbol: KLRC1

Gene Alias: CD159A, MGC13374, MGC59791, NKG2,

NKG2A

Gene Summary: Natural killer (NK) cells are

lymphocytes that can mediate lysis of certain tumor cells and virus-infected cells without previous activation. They can also regulate specific humoral and cell-mediated immunity. The protein encoded by this gene belongs to the killer cell lectin-like receptor family, also called NKG2 family, which is a group of transmembrane proteins preferentially expressed in NK cells. This family of proteins is characterized by the type II membrane orientation and the presence of a C-type lectin domain. This protein forms a complex with another family member, KLRD1/CD94, and has been implicated in the recognition of the MHC class I HLA-E molecules in NK cells. The genes of NKG2 family members form a killer cell lectin-like receptor gene cluster on chromosome 12. Four alternatively spliced transcript variants encoding two distinct isoforms have been observed. [provided by RefSeq]