



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

PTPRC (Human) Recombinant Protein (Biotin)

Catalog Number: P10484

Regulation Status: For research use only (RUO)

Product Description: Human PTPRC (P08575-4, Gln26-Lys416) partial recombinant protein with His-Avi tag at the C-Terminus expressed in HEK293 cells.

Sequence: Gln26-Lys416

Host: Human

Theoretical MW (kDa): 47.3

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Lyophilized

Conjugation: Biotin

Preparation Method: Mammalian cell (HEK293) expression system

Purity: > 95% by Tris-Bis PAGE
> 95% by HPLC

Endotoxin Level: < 0.1 EU per 1 ug as determined by the LAL method.

Activity: The EC₅₀ was 11.4 ng/mL, measured by ELISA at 1 ug/mL.

Recommend Usage: Biological Activity

ELISA

Tris-Bis PAGE

SEC-HPLC

The optimal working dilution should be determined by the end user.

Storage Buffer: Lyophilized from 0.22 um filtered solution in PBS, pH 7.4. (8% trehalose).

Storage Instruction: Store at -20°C for 12 months. After reconstitution, store at 4°C for 2-7 days, or store at -80°C for 3-6 months.

Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 5788

Gene Symbol: PTPRC

Gene Alias: B220, CD45, CD45R, GP180, LCA, LY5, T200

Gene Summary: The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP contains an extracellular domain, a single transmembrane segment and two tandem intracytoplasmic catalytic domains, and thus belongs to receptor type PTP. This gene is specifically expressed in hematopoietic cells. This PTP has been shown to be an essential regulator of T- and B-cell antigen receptor signaling. It functions through either direct interaction with components of the antigen receptor complexes, or by activating various Src family kinases required for the antigen receptor signaling. This PTP also suppresses JAK kinases, and thus functions as a regulator of cytokine receptor signaling. Four alternatively spliced transcripts variants of this gene, which encode distinct isoforms, have been reported. [provided by RefSeq]