

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

Datasheet

TNFRSF1B (Human) Recombinant Protein (Biotin)

Catalog Number: P10502

Regulation Status: For research use only (RUO)

Product Description: Human TNFRSF1B (P20333-1, Leu23-Asp257) partial recombinant protein with His tag

at the C-Terminus expressed in HEK293 cells.

Sequence: Leu23-Asp257

Host: Human

Theoretical MW (kDa): 26.2

Protocols: See our web site at

http://www.abnova.com/support/protocols.asp or product

page for detailed protocols

Form: Lyophilized

Conjugation: Biotin

Preparation Method: Mammalian cell (HEK293)

expression system

Purity: > 95% by Tris-Bis PAGE

> 95% by HPLC

Endotoxin Level: < 0.1 EU per 1 ug as determined by

the LAL method.

Activity: The EC₅₀ was 9.0 ng/mL, measured by ELISA

at 2 ug/mL.

Recommend Usage: Biological Activity

ELISA

Tris-Bis PAGE SEC-HPLC

The optimal working dilution should be determined by

the end user.

Storage Buffer: Lyophilized from 0.22 um filtered

solution in PBS, pH 7.4. (8% trehalose).

Storage Instruction: Store at -20°C for 12 months. After reconstitution, store at 4°C for 2-7 days, or store at

-80°C for 3-6 months.

Aliquot to avoid repeated freezing and thawing.

Entrez GenelD: 7133

Gene Symbol: TNFRSF1B

Gene Alias: CD120b, TBPII, TNF-R-II, TNF-R75, TNFBR, TNFR1B, TNFR2, TNFR80, p75, p75TNFR

Gene Summary: The protein encoded by this gene is a member of the TNF-receptor superfamily. This protein and TNF-receptor 1 form a heterocomplex that mediates the recruitment of two anti-apoptotic proteins, c-IAP1 and c-IAP2, which possess E3 ubiquitin ligase activity. The function of IAPs in TNF-receptor signalling is unknown, however, c-IAP1 is thought to potentiate TNFinduced apoptosis by the ubiquitination and degradation of TNF-receptor-associated factor 2, which mediates antiapoptotic signals. Knockout studies in mice also suggest a role of this protein in protecting neurons from apoptosis by stimulating antioxidative pathways. [provided by RefSeq]