



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

KDR (Human) Recombinant Protein

Catalog Number: P10564

Regulation Status: For research use only (RUO)

Product Description: Human KDR partial recombinant protein with His tag at the C-terminus expressed in HEK293 cells.

Sequence: Met1-Glu764

Host: Human

Theoretical MW (kDa): 87.08

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Lyophilized

Preparation Method: Mammalian cell (HEK293) expression system

Purification: Ni-NTA chromatography

Purity: > 95% as determined by SDS-PAGE.

Endotoxin Level: < 0.1 EU/ ug of protein by the LAL method.

Recommend Usage: SDS-PAGE

The optimal working dilution should be determined by the end user.

Storage Buffer: Lyophilized from a 0.2 um filtered solution of PBS, pH 7.4.

Storage Instruction: Store at -20°C for 12 months in lyophilized state.
After reconstitution with deionized water, store at -20 or -80°C for 1 month.
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 3791

Gene Symbol: KDR

Gene Alias: CD309, FLK1, VEGFR, VEGFR2

Gene Summary: Vascular endothelial growth factor (VEGF) is a major growth factor for endothelial cells. This gene encodes one of the two receptors of the VEGF. This receptor, known as kinase insert domain receptor, is a type III receptor tyrosine kinase. It functions as the main mediator of VEGF-induced endothelial proliferation, survival, migration, tubular morphogenesis and sprouting. The signalling and trafficking of this receptor are regulated by multiple factors, including Rab GTPase, P2Y purine nucleotide receptor, integrin α V β 3, T-cell protein tyrosine phosphatase, etc.. Mutations of this gene are implicated in infantile capillary hemangiomas. [provided by RefSeq]