



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

IFNG (Human) Recombinant Protein

Gene Alias: IFG, IFI

Catalog Number: P8159

Regulation Status: For research use only (RUO)

Product Description: Human IFNG (P01579, 24 a.a. - 161 a.a.) partial recombinant protein expressed in *Escherichia coli*.

Sequence:

MQDPYVKEAENLKKYFNAGHSDVADNGTLFLGILKNW
KEESDRKIMQSQIVSFYFKLFKNFKDDQSIQKSVETIKE
DMNVKFFNSNKKKRDDFEKLTNYSVTDLNVQRKAIHE
LIQVMAELSPAAGTKRKRSQMLFRG

Host: *Escherichia coli*

Theoretical MW (kDa): 16.3

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product
page for detailed protocols

Form: Liquid

Preparation Method: *Escherichia coli* expression
system

Purity: > 90% by SDS-PAGE

Activity: The ED₅₀ is < 0.5 ng/mL, measured in a
cytotoxicity assay using WiDr cells.

Recommend Usage: Biological Activity
SDS-PAGE

The optimal working dilution should be determined by
the end user.

Storage Buffer: In PBS pH 7.4 (10% glycerol)

Storage Instruction: Store at 2°C to 8°C for 1 week.
For long term storage, aliquot and store at -20°C to
-80°C.
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 3458

Gene Symbol: IFNG

Gene Summary: Interferon-gamma (IFNG), or type II interferon, is a cytokine critical for innate and adaptive immunity against viral and intracellular bacterial infections and for tumor control. Aberrant IFNG expression is associated with a number of autoimmune and inflammatory diseases. The importance of IFNG in the immune system stems in part from its ability to inhibit viral replication directly, but most importantly derives from its immunostimulatory and immunomodulatory effects. IFNG is produced predominantly by natural killer (NK) and natural killer T (NKT) cells as part of the innate immune response, and by CD4 (MIM 186940) and CD8 (see MIM 186910) cytotoxic T lymphocyte (CTL) effector T cells once antigen-specific immunity develops (Schoenborn and Wilson, 2007 [PubMed 17981204]).[supplied by OMIM]