



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

IL1B (Human) Recombinant Protein

Catalog Number: P8219

Regulation Status: For research use only (RUO)

Product Description: Human IL1B (P01584, 117 a.a. - 269 a.a.) partial recombinant protein expressed in HEK293 cells.

Sequence:

APVRSLNCTLRDSQQKSLVMSGPYELKALHLQGQDM
EQQVVFMSFVQGEESNDKIP VALGLKEKNLYLSCVL
KDDKPTLQLESVDPKNYPKKKMEKRFVFNKIEINNLE
FES AQFPNWIYSTQAENMPVFLGGTKGGQDITDFT
MQFVSS

Host: Mammals

Theoretical MW (kDa): 18-25

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product
page for detailed protocols

Form: Lyophilized

Preparation Method: Mammalian cell (HEK293)
expression system

Purity: > 95% by SDS-PAGE

Recommend Usage: Biological Activity
SDS-PAGE

The optimal working dilution should be determined by
the end user.

Storage Buffer: In sterile PBS containing 0.1%
endotoxin-free recombinant HSA.

Storage Instruction: Store at 2°C to 8°C for 1 week.
For long term storage, aliquot and store at -20°C to
-80°C.
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 3553

Gene Symbol: IL1B

Gene Alias: IL-1, IL1-BETA, IL1F2

Gene Summary: The protein encoded by this gene is a member of the interleukin 1 cytokine family. This cytokine is produced by activated macrophages as a proprotein, which is proteolytically processed to its active form by caspase 1 (CASP1/ICE). This cytokine is an important mediator of the inflammatory response, and is involved in a variety of cellular activities, including cell proliferation, differentiation, and apoptosis. The induction of cyclooxygenase-2 (PTGS2/COX2) by this cytokine in the central nervous system (CNS) is found to contribute to inflammatory pain hypersensitivity. This gene and eight other interleukin 1 family genes form a cytokine gene cluster on chromosome 2. [provided by RefSeq]