



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

IL5 (Human) Recombinant Protein

Catalog Number: P8278

Regulation Status: For research use only (RUO)

Product Description: Human IL5 (P05113, 20 a.a. - 134 a.a.) partial recombinant protein with His tag at C-terminus expressed in Sf9 cells.

Sequence:

ADPIPTSEIPTSALVKETLALLSTHRTLLIANETLRIPVPVH
KNHQLCTEEIFQGIGTLESQTVQGGTVERLFKNLSLIKK
YIDGQKKKCGEERRRVNQFLDYLQEFLGVMNTEWIIIE
SHHHHHH

Host: insect

Theoretical MW (kDa): 14.2

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product
page for detailed protocols

Form: Liquid

Preparation Method: Sf9 cell expression system

Purity: > 90% by SDS-PAGE

Activity: The ED₅₀ is < 1.5 ng/mL, measured in a cell
proliferation assay using TF-1 human erythroleukemic
cell.

Recommend Usage: Biological Activity

SDS-PAGE

The optimal working dilution should be determined by
the end user.

Storage Buffer: In PBS pH 7.4 (10% glycerol)

Storage Instruction: Store at 2°C to 8°C for 1 week.
For long term storage, aliquot and store at -20°C to
-80°C.
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 3567

Gene Symbol: IL5

Gene Alias: EDF, IL-5, TRF

Gene Summary: The protein encoded by this gene is a cytokine that acts as a growth and differentiation factor for both B cells and eosinophils. This cytokine is a main regulator of eosinopoiesis, eosinophil maturation and activation. The elevated production of this cytokine is reported to be related to asthma or hypereosinophilic syndromes. The receptor of this cytokine is a heterodimer, whose beta subunit is shared with the receptors for interleukine 3 (IL3) and colony stimulating factor 2 (CSF2/GM-CSF). This gene, together with those for interleukin 4 (IL4), interleukin 13 (IL13), and CSF2, form a cytokine gene cluster on chromosome 5. This cytokine, IL4, and IL13 are found to be regulated coordinately by long-range regulatory elements spread over 120 kilobases on chromosome 5q31. [provided by RefSeq]