



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

IL12B (Human) Recombinant Protein

Catalog Number: P8329

Regulation Status: For research use only (RUO)

Product Description: Human IL12B (P29460, 23 a.a. - 328 a.a.) partial recombinant protein with His tag at C-terminus expressed in Baculovirus.

Sequence:

ADPIWELKKDVYVVELDWYPDAPGEMVVLTCDTPEED
GITWTLQSSSEVLGSGKTLTIQVKEFGDAGQYTCHKG
GEVLSHSLLLLHKKEDGIWSTDILKDQKEPKNKTFRLC
EAKNYSGRFTCWWLTTISTDLTFSVKSSRGSSDPQGV
TCGAATLSAERVGRDNKEYEYSVEQEDSACPAAEE
SLPIEVMVDAVHKLKYENYTSSFFIRDIIKPDPPKNLQLK
PLKNSRQVEVSWEYPDTWSTPHSYFSLTFCVQVQGK
SKREKKDRVFTDKTSATVICRKNASISVRAQDRYYSSS
WSEWASVPCSHHHHHH

Host: Viruses

Theoretical MW (kDa): 35.8

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product
page for detailed protocols

Form: Liquid

Preparation Method: Baculovirus

Purity: > 90% by SDS-PAGE

Recommend Usage: Biological Activity
SDS-PAGE

The optimal working dilution should be determined by
the end user.

Storage Buffer: In 20mM Tris-HCl buffer pH 8.0
(100mM NaCl, 2mM DTT, 100mM PMSF, 20% glycerol)

Storage Instruction: Store at 2°C to 8°C for 1 week.
For long term storage, aliquot and store at -20°C to
-80°C.
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 3593

Gene Symbol: IL12B

Gene Alias: CLMF, CLMF2, IL-12B, NKSF, NKSF2

Gene Summary: This gene encodes a subunit of interleukin 12, a cytokine that acts on T and natural killer cells, and has a broad array of biological activities. Interleukin 12 is a disulfide-linked heterodimer composed of the 40 kD cytokine receptor like subunit encoded by this gene, and a 35 kD subunit encoded by IL12A. This cytokine is expressed by activated macrophages that serve as an essential inducer of Th1 cells development. This cytokine has been found to be important for sustaining a sufficient number of memory/effector Th1 cells to mediate long-term protection to an intracellular pathogen. Overexpression of this gene was observed in the central nervous system of patients with multiple sclerosis (MS), suggesting a role of this cytokine in the pathogenesis of the disease. The promoter polymorphism of this gene has been reported to be associated with the severity of atopic and non-atopic asthma in children. [provided by RefSeq]