

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com



9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

Datasheet

FGF8 (Human) Recombinant Protein

Catalog Number: P8606

Regulation Status: For research use only (RUO)

Product Description: Human FGF8 recombinant protein expressed in?HEK293 cells.

Host: Human

Theoretical MW (kDa): 30-45

Protocols: See our web site at http://www.abnova.com/support/protocols.asp or product page for detailed protocols

Form: Lyophilized

Preparation Method: Mammalian cell (HEK 293) expression system

Purification: chromatography

Purity: > 95% as determined by SDS-PAGE.

Activity: $ED_{50} < 60$ ng/mL, determined by dosedependent stimulation of the proliferation of the Balb/3T3 cell line.

Storage Buffer: Lyophilized from a solution containing 10mM Tris-HCl, pH 7.4, 800mM NaCl. Reconstitute the lyophilized powder in ddH_2O containing 0.1% endotoxin-free recombinant HSA. to 100 ug/mL.

Storage Instruction: Lyophilized protein at room temperature for 3 weeks, should be stored at -20°C. Protein aliquots at 4°C for 2-7 days and should be stored at -20°C to -80°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).

Avoid repeated freeze/thaw cycles.

Entrez GenelD: 2253

Gene Symbol: FGF8

Gene Alias: AIGF, HBGF-8, MGC149376

Gene Summary: The protein encoded by this gene is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. This protein is known to be a factor that supports androgen and anchorage independent growth of mammary tumor cells. Overexpression of this gene has been shown to increase tumor growth and angiogensis. The adult expression of this gene is restricted to testes and ovaries. Temporal and spatial pattern of this gene expression suggests its function as an embryonic epithelial factor. Studies of the mouse and chick homologs revealed roles in midbrain and limb development, organogenesis, embryo gastrulation and left-right axis determination. The alternative splicing of this gene results in four transcript variants. [provided by RefSeq]