



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

GDF11 (Human) Recombinant Protein

Catalog Number: P8698

Regulation Status: For research use only (RUO)

Product Description: Human GDF11 partial recombinant protein with His tag in N-terminus expressed in *Escherichia coli*.

Sequence:

MGSSHHHHHSSGLVPRGSHMGSNLGLDCDEHSSE
SRCCRYPLTVDFEAFGWDWIIAPKRYKANYCSGQCEY
MFMQYPHTHLVQQANPRGSAGPCCTPTKMSPINML
YFNDKQQIYGKIPGMVVDRCGCS

Host: *Escherichia coli*

Theoretical MW (kDa): 14.8

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Preparation Method: *Escherichia coli* expression system

Purification: chromatographic

Purity: > 85% as determined by SDS-PAGE.

Storage Buffer: Solution (1 mg/mL) containing 20 mM Tris-HCl, pH 8.0, 10% glycerol.

Storage Instruction: Store at 4°C for 2-4 weeks and should be stored at -20°C to -80°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).
Avoid repeated freeze/thaw cycles.

Entrez GeneID: 10220

Gene Symbol: GDF11

Gene Alias: BMP-11, BMP11

Gene Summary: The protein encoded by this gene is a

member of the bone morphogenetic protein (BMP) family and the TGF-beta superfamily. This group of proteins is characterized by a polybasic proteolytic processing site which is cleaved to produce a mature protein containing seven conserved cysteine residues. The members of this family are regulators of cell growth and differentiation in both embryonic and adult tissues. Studies in mice and *Xenopus* suggest that this protein is involved in mesodermal formation and neurogenesis during embryonic development. [provided by RefSeq]