



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

IL28A (Human) Recombinant protein

Catalog Number: P8815

Regulation Status: For research use only (RUO)

Product Description: Human IL28A (Q8IZJ0, 26 a.a. - 200 a.a) partial recombinant protein with His tag at N-terminal expressed in *Escherichia Coli*.

Sequence: MGSSHHHHHH SSGLVPRGSH

MGSPVARLH	GALPDARGCH	IAQFKSLSPQ
ELQAFKRAKD	ALEESLLLKD	CRCHSRLFPR
TWDLRQLQVR	ERPMALAEAL	ALTLKVLEAT
ADTDPALVDV	LDQPLHTLHH	ILSQFRACIQ
PQPTAGPRTR	GRLHHWLYRL	QEAPKKESPG
CLEASVTFNL	FRLLTRDLNC	VASGDLCV

Host: *Escherichia coli*

Theoretical MW (kDa): 22.1

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product
page for detailed protocols

Form: Liquid

Preparation Method: *Escherichia coli* expression
system

Purity: > 85% by SDS-PAGE.

Storage Buffer: In 20mM Tris-HCl, pH 8.0, 50% glycerol
and 0.4M Urea

Storage Instruction: Store at 4°C for 2~4 week. For
long term storage store at -20°C. For long term storage it
is recommended to add a carrier protein (0.1% HSA or
BSA).
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 282616

Gene Symbol: IL28A

Gene Alias: IFNL2, IL-28A

Gene Summary: This gene encodes a cytokine distantly
related to type I interferons and the IL-10 family. This

gene, interleukin 28B (IL28B), and interleukin 29 (IL29)
are three closely related cytokine genes that form a
cytokine gene cluster on a chromosomal region mapped
to 19q13. Expression of the cytokines encoded by the
three genes can be induced by viral infection. All three
cytokines have been shown to interact with a
heterodimeric class II cytokine receptor that consists of
interleukin 10 receptor, beta (IL10RB) and interleukin 28
receptor, alpha (IL28RA). [provided by RefSeq]