



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

IL29 (Human) Recombinant protein

Catalog Number: P8820

Regulation Status: For research use only (RUO)

Product Description: Human IL29 (Q8IU54, 20 a.a. - 200 a.a) partial recombinant protein with His tag at C-terminal expressed in Sf9 cells.

Sequence: GPVPTSKPTT TGKGCHIGRF

KSLSPQELAS	FKKARDALEE	SLKLKNWSCS
SPVFPGNWDL	RLLQVRERP	ALEAELALTL
KVLEAAAGPA	LEDVLDQPLH	TLHHILSQLQ
ACIQPQPTAG	PRPRGRLHHW	LHRLQEAPKK
ESAGCLEASV	TFNLFRLLTR	DLKYVADGNL
CLRTSTHPES	THHHHHH	

Host: insect

Theoretical MW (kDa): 20.8

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product
page for detailed protocols

Form: Liquid

Preparation Method: Insect cell (Sf9) expression
system

Purity: > 90% by SDS-PAGE.

Storage Buffer: In PBS, pH 7.4 and 10% glycerol

Storage Instruction: Store at 4°C for 2~4 week. For
long term storage store at -20°C. For long term storage it
is recommended to add a carrier protein (0.1% HSA or
BSA).
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 282618

Gene Symbol: IL29

Gene Alias: IFNL1, IL-29

Gene Summary: This gene encodes a cytokine distantly
related to type I interferons and the IL-10 family. This
gene, interleukin 28A (IL28A), and interleukin 28B

(IL28B) are three closely related cytokine genes that
form a cytokine gene cluster on a chromosomal region
mapped to 19q13. Expression of the cytokines encoded
by the three genes can be induced by viral infection. All
three cytokines have been shown to interact with a
heterodimeric class II cytokine receptor that consists of
interleukin 10 receptor, beta (IL10RB) and interleukin 28
receptor, alpha (IL28RA). [provided by RefSeq]