



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

TNFRSF1B (Human) Recombinant Protein

Catalog Number: P9246

Regulation Status: For research use only (RUO)

Product Description: Human TNFRSF1B partial recombinant protein with His tag in C-terminus expressed in Baculovirus cells.

Sequence:

LPAQVAFTPYAPEPGSTCRLREYYDQTAQMCCSKCS
PGQHAKVFCTKTSDTVCDSCEDSTYQLWNWVPECL
SCGSRSSDQVETQACTREQNRICRPGWYCALSK
QEGCRLCAPLRKCRPGFGVARPGTETSDVVCPCAP
GTFSNTTSSTDICRPHQICNVVAIPGNASMDAVCTSTS
PTRSMAPGAVHLPQPVSTRSQHTQPTPEPSTAPSTSF
LLPMGPPPAEGSTGDHHHHHH

Host: Viruses

Theoretical MW (kDa): 25.9

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product
page for detailed protocols

Form: Liquid

Preparation Method: Baculovirus expression system

Purification: chromatographic

Purity: > 95% as determined by SDS-PAGE.

Storage Buffer: Solution (1 mg/mL) containing 1X PBS,
pH 7.4, 10% glycerol.

Storage Instruction: Store at 4°C for one weeks and
should be stored at -20°C to -80°C. For long term
storage it is recommended to add a carrier protein (0.1%
HSA or BSA).
Avoid repeated freeze/thaw cycles.

Entrez GeneID: 7133

Gene Symbol: TNFRSF1B

Gene Alias: CD120b, TBPII, TNF-R-II, TNF-R75,

TNFR, TNFR1B, TNFR2, TNFR80, p75, p75TNFR

Gene Summary: The protein encoded by this gene is a member of the TNF-receptor superfamily. This protein and TNF-receptor 1 form a heterocomplex that mediates the recruitment of two anti-apoptotic proteins, c-IAP1 and c-IAP2, which possess E3 ubiquitin ligase activity. The function of IAPs in TNF-receptor signalling is unknown, however, c-IAP1 is thought to potentiate TNF-induced apoptosis by the ubiquitination and degradation of TNF-receptor-associated factor 2, which mediates anti-apoptotic signals. Knockout studies in mice also suggest a role of this protein in protecting neurons from apoptosis by stimulating antioxidative pathways. [provided by RefSeq]