



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

TNFRSF8 (Human) Recombinant Protein

Catalog Number: P9257

Regulation Status: For research use only (RUO)

Product Description: Human TNFRSF8 partial recombinant protein with His tag in C-terminus expressed in Baculovirus cells.

Sequence:

ADPFPQDRPFEDTCHGNPSHYDYKAVRRCCYRCPMG
LFPTQQCPQRPTDCRKQCEPDYYLDEADRCTACVTC
SRDDLVEKTPCAWNSSRVCECRPGMFCSTSAVNCA
RCFFHSVCPAGMIVKFGTAQKNTVCEPASPGVSPAC
ASPENCKEPSSGTIPQAKPTPVSPATSSASTMPVRGG
TRLAQEAASKLTRAPDSPSSVGRPSSDPGLSPTQPCP
EGSGDCRKQCEPDYYLDEAGRCTACVSCSRDDLVEK
TPCAWNSSRTCECRPGMICATSATNSCARCVPYPICA
AETVTKPQDMAEKDTTFEAPPLGTQPD CNPTPENGEA
PASTSPTQSLLVDSQASKTLPIPTSAPVALSSTGKHHH
HHH

Host: Viruses

Theoretical MW (kDa): 39.5

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product
page for detailed protocols

Form: Liquid

Preparation Method: Baculovirus expression system

Purification: chromatographic

Purity: > 95% as determined by SDS-PAGE.

Storage Buffer: Solution (1 mg/mL) containing 1X PBS,
pH 7.4, 10% glycerol.

Storage Instruction: Store at 4°C for one weeks and
should be stored at -20°C to -80°C. For long term
storage it is recommended to add a carrier protein (0.1%
HSA or BSA).

Avoid repeated freeze/thaw cycles.

Entrez GeneID: 943

Gene Symbol: TNFRSF8

Gene Alias: CD30, D1S166E, KI-1

Gene Summary: The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is expressed by activated, but not by resting, T and B cells. TRAF2 and TRAF5 can interact with this receptor, and mediate the signal transduction that leads to the activation of NF-kappaB. This receptor is a positive regulator of apoptosis, and also has been shown to limit the proliferative potential of autoreactive CD8 effector T cells and protect the body against autoimmunity. Two alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported. [provided by RefSeq]