



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

VEGFA (Human) Recombinant Protein

Catalog Number: P9307

Regulation Status: For research use only (RUO)

Product Description: Human VEGFA recombinant protein expressed in Insect Cells.

Sequence:

APMAEGGGQNHHEVVKFMDVYQRSYCHPIETLVDIFQ
EYPDEIEYIFKPSCVPLMRGCGCCNDEGLECVPTES
NITMQIMRIKPHQGQHIGEMSFLQHNKCECRPKKDRA
RQEKCDKPRR

Host: insect

Theoretical MW (kDa): 36

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product
page for detailed protocols

Form: Lyophilized

Preparation Method: Insect cells expression system

Purification: chromatographic

Purity: > 95% as determined by SDS-PAGE.

Activity: ED₅₀ is 2-10 ng/mL, measured in a cell
proliferation assay using primary HUVEC cells.

Storage Buffer: Lyophilized from a solution containing
50 mM acetic acid. Reconstitute the lyophilized powder
in 50 mM acetic acid to 50 ug/mL.

Storage Instruction: Lyophilized protein at room
temperature for 3 weeks, should be stored at -20°C.
Protein aliquots at 4°C for 2-7 days and should be stored
at -20°C to -80°C. For long term storage it is
recommended to add a carrier protein (0.1% HSA or
BSA).
Avoid repeated freeze/thaw cycles.

Entrez GeneID: 7422

Gene Symbol: VEGFA

Gene Alias: MGC70609, VEGF, VEGF-A, VPF

Gene Summary: This gene is a member of the PDGF/VEGF growth factor family and encodes a protein that is often found as a disulfide linked homodimer. This protein is a glycosylated mitogen that specifically acts on endothelial cells and has various effects, including mediating increased vascular permeability, inducing angiogenesis, vasculogenesis and endothelial cell growth, promoting cell migration, and inhibiting apoptosis. Elevated levels of this protein is linked to POEMS syndrome, also known as Crow-Fukase syndrome. Mutations in this gene have been associated with proliferative and nonproliferative diabetic retinopathy. Alternatively spliced transcript variants, encoding either freely secreted or cell-associated isoforms, have been characterized. There is also evidence for the use of non-AUG (CUG) translation initiation sites upstream of, and in-frame with the first AUG, leading to additional isoforms. [provided by RefSeq]