



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

CCL2 (Human) Recombinant Protein

Catalog Number: P9508

Regulation Status: For research use only (RUO)

Product Description: Human CCL2 (P13500, 24 a.a. - 99 a.a.) partial recombinant protein with His tag at C-terminus expressed in HEK293 cells.

Sequence:

QPDAINAPVTCCYNFTNRKISVQRLASYRRITSSKCPK
EAVIFKTIVAKEICADPKQKWVQDSMDHLDKQTQTPKT
HHHHHH

Host: Mammals

Theoretical MW (kDa): 9.5

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product
page for detailed protocols

Form: Liquid

Preparation Method: Mammalian cell (HEK293)
expression system

Purity: > 90.0% by SDS-PAGE

Recommend Usage: Biological Activity
SDS-PAGE
The optimal working dilution should be determined by
the end user.

Storage Buffer: In PBS pH 7.4 (10% glycerol)

Storage Instruction: Store at 2°C to 8°C for 1 week.
For long term storage, aliquot and store at -20°C to
-80°C.
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 6347

Gene Symbol: CCL2

Gene Alias: GDCE-2, HC11, HSMCR30, MCAF,
MCP-1, MCP1, MGC9434, SCYA2, SMC-CF

Gene Summary: This gene is one of several cytokine

genes clustered on the q-arm of chromosome 17. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The protein encoded by this gene is structurally related to the CXC subfamily of cytokines. Members of this subfamily are characterized by two cysteines separated by a single amino acid. This cytokine displays chemotactic activity for monocytes and basophils but not for neutrophils or eosinophils. It has been implicated in the pathogenesis of diseases characterized by monocytic infiltrates, like psoriasis, rheumatoid arthritis and atherosclerosis. It binds to chemokine receptors CCR2 and CCR4. [provided by RefSeq]