



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

CCL13 (Human) Recombinant Protein

Catalog Number: P9519

Regulation Status: For research use only (RUO)

Product Description: Human CCL13 (Q99616, 24 a.a. - 98 a.a.) partial recombinant protein expressed in *Escherichia coli*.

Sequence:

QPDALNVPSTCCFTFSSKKISLQRLKSYVITTSRCPQK
AVIFRTKLGKEICADPKEKWVQNYMKHLGRKAHTLKT

Host: *Escherichia coli*

Theoretical MW (kDa): 8.6

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product
page for detailed protocols

Form: Lyophilized

Preparation Method: *Escherichia coli* expression
system

Purity: > 96.0% by SDS-PAGE

Activity: The specific activity as determined by the
ability of MCP-4 to chemoattract human monocytes at 10
- 100 ng/mL, corresponding to a specific activity of
10000 - 100000 units/mg.

Recommend Usage: Biological Activity
SDS-PAGE

The optimal working dilution should be determined by
the end user.

Storage Buffer: Lyophilized from sterile distilled Water
is > 100 ug/mL

Storage Instruction: Store at 2°C to 8°C for 1 week.
For long term storage, aliquot and store at -20°C to
-80°C.
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 6357

Gene Symbol: CCL13

Gene Alias: CKb10, MCP-4, MGC17134, NCC-1,
NCC1, SCYA13, SCYL1

Gene Summary: This gene is one of several Cys-Cys
(CC) cytokine genes clustered on the q-arm of
chromosome 17. Cytokines are a family of secreted
proteins involved in immunoregulatory and inflammatory
processes. The CC cytokines are proteins characterized
by two adjacent cysteines. The cytokine encoded by this
gene displays chemotactic activity for monocytes,
lymphocytes, basophils and eosinophils, but not
neutrophils. This chemokine plays a role in accumulation
of leukocytes during inflammation. It may also be
involved in the recruitment of monocytes into the arterial
wall during atherosclerosis. [provided by RefSeq]