



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

CCL18 (Human) Recombinant Protein

Catalog Number: P9550

Regulation Status: For research use only (RUO)

Product Description: Human CCL18 (P55774, 21 a.a. - 89 a.a.) partial recombinant protein expressed in *Escherichia coli*.

Sequence:

AQVGTKNELCCLVYTSWQIPQKFIVDYSETSPQCPKP
GVILLTKRGRQICADPNKKWVQKYISDLKLNA

Host: *Escherichia coli*

Theoretical MW (kDa): 7.8

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product
page for detailed protocols

Form: Lyophilized

Preparation Method: *Escherichia coli* expression
system

Purity: > 97.0% by SDS-PAGE

Activity: The Biological activity is 1 - 10 ng/mL, was
calculated by the ability to chemoattract primary Human
T lymphocytes, corresponding to a specific activity of
100000 - 1000000 IU/mg.

Recommend Usage: Biological Activity
SDS-PAGE

The optimal working dilution should be determined by
the end user.

Storage Buffer: Lyophilized from sterile distilled Water
is > 100 ug/mL

Storage Instruction: Store at 2°C to 8°C for 1 week.
For long term storage, aliquot and store at -20°C to
-80°C.
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 6362

Gene Symbol: CCL18

Gene Alias: AMAC-1, AMAC1, CKb7, DC-CK1, DCCK1,
MIP-4, PARC, SCYA18

Gene Summary: This gene is one of several Cys-Cys
(CC) cytokine genes clustered on the q arm of
chromosome 17. Cytokines are a family of secreted
proteins involved in immunoregulatory and inflammatory
processes. The CC cytokines are proteins characterized
by two adjacent cysteines. The cytokine encoded by this
gene displays chemotactic activity for naive T cells,
CD4+ and CD8+ T cells and nonactivated lymphocytes,
but not for monocytes or granulocytes. This chemokine
attracts naive T lymphocytes toward dendritic cells and
activated macrophages in lymph nodes. It may play a
role in both humoral and cell-mediated immunity
responses. [provided by RefSeq]