



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

CDKN1A (Human) Recombinant Protein

Catalog Number: P9610

Regulation Status: For research use only (RUO)

Product Description: Human CDKN1A (P38936, 1 a.a. - 164 a.a.) full recombinant protein with His tag at N-terminus expressed in *Escherichia coli*.

Sequence:

MGSSHHHHHSSGLVPRGSHMSEPAGDVRQNPCGS
KACRRLFPGPVDSEQLSRDCDALMAGCIQEARERWNF
DFVTETPLEGDFAWERVRLGLPKLYLPTGPRRGRDE
LGGRRPGTSPALLQGTAEEDHVDLSLSCTLVPRSGE
QAEGLSPGGPGDSQGRKRRQTSMTDFYHSKRRLIFSK
RKP

Host: *Escherichia coli*

Theoretical MW (kDa): 20.2

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product
page for detailed protocols

Form: Liquid

Preparation Method: *Escherichia coli* expression
system

Purity: > 85.0% by SDS-PAGE

Recommend Usage: Biological Activity

SDS-PAGE

The optimal working dilution should be determined by
the end user.

Storage Buffer: In 20mM Tris-HCl pH 8.0 (2 M Urea,
0.1 M NaCl and 10% glycerol)

Storage Instruction: Store at 2°C to 8°C for 1 week.
For long term storage, aliquot and store at -20°C to
-80°C.

Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 1026

Gene Symbol: CDKN1A

Gene Alias: CAP20, CDKN1, CIP1, MDA-6, P21, SDI1,
WAF1, p21CIP1

Gene Summary: This gene encodes a potent cyclin-dependent kinase inhibitor. The encoded protein binds to and inhibits the activity of cyclin-CDK2 or -CDK4 complexes, and thus functions as a regulator of cell cycle progression at G1. The expression of this gene is tightly controlled by the tumor suppressor protein p53, through which this protein mediates the p53-dependent cell cycle G1 phase arrest in response to a variety of stress stimuli. This protein can interact with proliferating cell nuclear antigen (PCNA), a DNA polymerase accessory factor, and plays a regulatory role in S phase DNA replication and DNA damage repair. This protein was reported to be specifically cleaved by CASP3-like caspases, which thus leads to a dramatic activation of CDK2, and may be instrumental in the execution of apoptosis following caspase activation. Two alternatively spliced variants, which encode an identical protein, have been reported. [provided by RefSeq]