

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



## Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

#### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

### **Datasheet**

#### **IL4 (Human) Recombinant Protein**

Catalog Number: P9954

Regulation Status: RUO grade

**Product Description:** Human IL4 recombinant protein with His tag at the C-terminus expressed in *Escherichia* 

coli.

Host: Escherichia coli

Theoretical MW (kDa): Calculated MW: 15.9

Regulatory Status: RUO grade

Protocols: See our web site at

http://www.abnova.com/support/protocols.asp or product

page for detailed protocols

Form: Lyophilized

Preparation Method: Escherichia coli expression

system

Purity: > 98% by SDS-PAGE

Endotoxin Level: < 0.1 EU/ug

**Activity:** The ED<sub>50</sub> for this effect is < 0.2 ng/mL, measured by its ability to induce TF-1 cells proliferation. The specific activity of recombinant human IL-4 is approximately > 1 x  $10^7$  IU/mg.

Recommend Usage: SDS-PAGE

The optimal working dilution should be determined by

the end user.

**Storage Buffer:** Lyophilized from PBS, pH 8.0. Reconstitute the lyophilized protein in sterile  $H_2O$  to a concentration of at least 200 ug/mL and incubate the stock solution for at least 20 min to ensure sufficient redissolution. Please use the protein within one month after reconstitution.

Storage Instruction: Store at -20°C for 12 months in

lyophilized state.

After reconstitution with deionized water, store at -20 or -80°C for 1 month.

Aliquot to avoid repeated freezing and thawing.

Entrez GenelD: 3565

Gene Symbol: IL4

Gene Alias: BCGF-1, BCGF1, BSF1, IL-4, MGC79402

Gene Summary: The protein encoded by this gene is a pleiotropic cytokine produced by activated T cells. This cytokine is a ligand for interleukin 4 receptor. The interleukin 4 receptor also binds to IL13, which may contribute to many overlapping functions of this cytokine and IL13. STAT6, a signal transducer and activator of transcription, has been shown to play a central role in mediating the immune regulatory signal of this cytokine. This gene, IL3, IL5, IL13, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL13. This gene, IL13 and IL5 are found to be regulated coordinately by several long-range regulatory elements in an over 120 kilobase range on the chromosome. Two alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported. [provided by RefSeq]