



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



Datasheet

IL13RA1 (Human) Recombinant Protein

Catalog Number: P9965

Regulation Status: For research use only (RUO)

Product Description: Human IL13RA1 (P78552-1, Ala27-Thr343) partial recombinant protein with His tag at C-terminus expressed in HEK293 cells.

Sequence: Ala27-Thr343

Host: Human

Theoretical MW (kDa): 37.7 kDa

Due to g

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Preparation Method: Mammalian cell (HEK293) expression system

Purity: > 95% as determined by Tris-Bis PAGE; > 95% as determined by HPLC

Endotoxin Level: < 1 EU per 1 ug of protein
(determined by LAL method)

Recommend Usage: Biological Activity

SPR

Tris-Bis PAGE

The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS, pH 7.4.

Storage Instruction: Store at -80°C for 1 year.
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 3597

Gene Symbol: IL13RA1

Gene Alias: CD213A1, IL-13Ra, NR4

Gene Summary: The protein encoded by this gene is a subunit of the interleukin 13 receptor. This subunit forms a receptor complex with IL4 receptor alpha, a subunit shared by IL13 and IL4 receptors. This subunit serves as a primary IL13-binding subunit of the IL13 receptor, and may also be a component of IL4 receptors. This protein has been shown to bind tyrosine kinase TYK2, and thus may mediate the signaling processes that lead to the activation of JAK1, STAT3 and STAT6 induced by IL13 and IL4. [provided by RefSeq]