



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

FLT4 (Human) Recombinant Protein

Catalog Number: P9987

Regulation Status: For research use only (RUO)

Product Description: Human FLT4 (P35916-1, Tyr25-Ile776) partial recombinant protein with His-Avi tag at C-terminus expressed in HEK293 cells.

Sequence: Tyr25-Ile776

Host: Human

Theoretical MW (kDa): 87.4 kDa

Due to g

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product
page for detailed protocols

Form: Lyophilized

Preparation Method: Mammalian cell (HEK293)
expression system

Purity: > 95% as determined by Tris-Bis PAGE; > 95%
as determined by HPLC

Endotoxin Level: < 1 EU per 1 ug of protein
(determined by LAL method)

Recommend Usage: Biological Activity

SPR

Tris-Bis PAGE

The optimal working dilution should be determined by
the end user.

Storage Buffer: Lyophilized from filtered solution in
PBS, pH 7.4 (8% trehalose).

Storage Instruction: After reconstitution with deionized
water to a final concentration more than 100 ug/ml, store
at 4°C for 1 week. For long term storage, store at -80°C
for 1 year.

Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 2324

Gene Symbol: FLT4

Gene Alias: FLT41, LMPH1A, PCL, VEGFR3

Gene Summary: This gene encodes a tyrosine kinase
receptor for vascular endothelial growth factors C and D.
The protein is thought to be involved in
lymphangiogenesis and maintenance of the lymphatic
endothelium. Mutations in this gene cause hereditary
lymphedema type IA. [provided by RefSeq]