



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

CCDC50 polyclonal antibody

Catalog Number: PAB20051

Regulatory Status: For research use only (RUO)

Product Description: Rabbit polyclonal antibody raised against recombinant CCDC50.

Immunogen: Recombinant protein corresponding to amino acids of human CCDC50.

Sequence:

QEKKDEDIARLLQEKELQEEKRKKHFPEFPATRAYAD
SYYEDGGMKPRVMKEAVSTPSRMAHRDQEWYDAEI
ARKLQEEELLATQVDMRAAQVAQDEEIARLLMAEEKK
AYKKAKEREKSSLDKQDPEWKPKTAKAANSKSKES
DE

Host: Rabbit

Reactivity: Human, Mouse, Rat

Applications: IF, IHC-P, WB, WB-Ce
(See our web site product page for detailed applications information)

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Purification: Antigen affinity purification

Isotype: IgG

Recommend Usage: Immunohistochemistry
(1:1000-1:2500)
Western Blot (1:250-1:500)
Immunofluorescence (1-4 ug/mL)
The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS, pH 7.2 (40% glycerol, 0.02% sodium azide)

Storage Instruction: Store at 4°C. For long term storage store at -20°C.
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 152137

Gene Symbol: CCDC50

Gene Alias: C3orf6, DFNA44, YMER

Gene Summary: This gene encodes a soluble, cytoplasmic, tyrosine-phosphorylated protein with multiple ubiquitin-interacting domains. Mutations in this gene cause nonsyndromic, postlingual, progressive sensorineural DFNA44 hearing loss. In mouse, the protein is expressed in the inner ear during development and postnatal maturation and associates with microtubule-based structures. This protein may also function as a negative regulator of NF- κ B signaling and as an effector of epidermal growth factor (EGF)-mediated cell signaling. Alternative splicing results in multiple transcript variants encoding distinct isoforms. [provided by RefSeq]