



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

MARCO polyclonal antibody

Catalog Number: PAB31836

Regulatory Status: For research use only (RUO)

Product Description: Rabbit polyclonal antibody raised against recombinant human MARCO.

Sequence:

LNLQARLRVLEMYFLNDTLAAEDSPSFSLLQSAHPGE
HLAQGASRLQVLQAQLTWVRV

Host: Rabbit

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product
page for detailed protocols

Specificity: This antibody reacts to MARCO.

Form: Liquid

Purification: Affinity purification

Isotype: IgG

Recommend Usage: Immunofluorescence (0.25-2
ug/mL)
Immunohistochemistry (1:500 - 1:1000)
The optimal working dilution should be determined by
the end user.

Storage Buffer: In PBS, pH7.2 (40% glycerol, 0.02%
sodium azide).

Storage Instruction: Store at 4°C. For long term
storage store at -20°C.
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 8685

Gene Symbol: MARCO

Gene Alias: SCARA2

Gene Summary: The protein encoded by this gene is a
member of the class A scavenger receptor family and is
part of the innate antimicrobial immune system. The
protein may bind both Gram-negative and Gram-positive

bacteria via an extracellular, C-terminal, scavenger
receptor cysteine-rich (SRCR) domain. In addition to
short cytoplasmic and transmembrane domains, there is
an extracellular spacer domain and a long, extracellular
collagenous domain. The protein may form a trimeric
molecule by the association of the collagenous domains
of three identical polypeptide chains. [provided by
RefSeq]