



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

For research use only
Not for use in diagnostic procedures

iMatrix-332

Product No. AMS.892 031 350 µg
Product No. AMS.892 032 1,050 µg

Version 002
Store at 2-15°C

Product description: iMatrix-332 is a recombinant human laminin-332 E8 fragment protein expressed in Chinese Hamster Ovary (CHO)-S cells. iMatrix-332 contains the integrin-binding site of the laminin-332 molecule. iMatrix-332 is useful for differentiation of iPSC and ESCs into corneal epithelial cells. iMatrix-332 also supports the culture of the other cells adhering to laminin-332 such as keratinocytes.

Content: Recombinant human laminin-332 E8 fragment protein in 20 mM phosphate buffer, 500 mM NaCl

Protein concentration: 0.5 mg/mL

Amount: 175 µg / 0.35 mL / tube
Product No. AMS.892 031 350 µg / 2 tubes
Product No. AMS.892 032 1,050 µg / 6 tubes

Storage: Store at 2°C to 15°C, protect from light.

Expiration date: The shelf life is 2 years from the date of manufacture. The expiration date is printed on the carton.

Activity: The dissociation constant (Kd) for the binding with integrin α3β1 is 5 nM or less.

Methods of use: By the following method iMatrix-332 can be coated onto a culture vessel. The optimum coating density may differ by cell-type, cell-line, medium selected, or purpose. Insufficient coating density may result in the detachment of cells and varied cell conditions while the excessive coating density may lead to difficulty in detaching cells for passage.

Coating protocol

Determine the optimal coating density. 0.5 µg/cm² is a standard but test between 0.1 and 1.5 µg/cm².

- 1) Dilute iMatrix-332 with PBS(-). Use the diluted iMatrix-332 immediately. To coat with 0.5 µg/cm² onto a 6-well plate with 9.6 cm²/well, dilute 9.6 µL of iMatrix-332 with 2 mL of PBS(-) per well.
- 2) Place the diluted iMatrix-332 into a culture vessel and incubate either at 37°C for 1 h, or at room temperature for 3 h, or at 4°C overnight.
- 3) Aspirate the coating solution. Then, immediately seed your cells. Do not allow the coated surface to dry.

*If you face difficulties in detaching cells for passage, re-adjust the conditions (e.g., reduce the coating density).

*Coating protocol is illustrative only.

References:

Nishiuchi R. *et al.* (2006) *Matrix Biol.* **25** (3), 189-97
Miyazaki T. *et al.* (2012) *Nat. Commun.* **3**, 1236
Shibata S. *et al.* (2018) *Cell Rep.* **25** (6), 1668-79
Shibata S. *et al.* (2020) *Stem Cell Reports* **14** (4), 663-76

Caution: For research use only. Not intended for human use. In the event of accidental ingestion or contact with the eyes, immediately wash the affected area and seek medical attention.

Product information: Current information including references and Q&A are available see <https://www.amsbio.com/imatrix-recombinant-laminin-series/>

Designed by: MATRIXOME, Inc.
3-2 Yamadaoka, Suita, Osaka 565-0871, Japan
Institute for Protein Research, Osaka University

Manufactured by: Nippi, Incorporated
1-1-1 Senju Midori-cho, Adachi, Tokyo 120-8601, Japan

AMSBIO | www.amsbio.com | info@amsbio.com

AMSBIO LLC
USA & Canada 
1035 Cambridge Street,
Cambridge, MA 02141
T: +1 (617) 945-5033 or
T: +1 (800) 987-0985
F: +1 (617) 945-8218

AMSBIO Europe BV
EU 
Berenkoog 41,
1822 BH Alkmaar,
Netherlands
T: +31 (0) 72 8080244
F: +31 (0) 72 8080142

AMS Biotechnology (Europe) Ltd
UK & Rest of the World 
184 Park Drive, Milton Park
Abingdon OX14 4SE
T: +44 (0) 1235 828 200
F: +44 (0) 1235 820 482

AMS Biotechnology (Europe) Ltd
Switzerland 
Via Lisano 3,
(CP.683)
CH-6900 Massagno
T: +41 (0) 91 604 55 22
F: +41 (0) 91 605 17 85

AMSBIO Europe BV
Deutschland 
T: +49 (0) 69 779099
F: +49 (0) 69 13376880